

RATE OF ESCAPE OF RANDOM WALKS ON FREE PRODUCTS

LORENZ A. GILCH

(February 1, 2008)

Abstract

Suppose we are given the free product V of a finite family of finite or countable sets $(V_i)_{i \in \mathcal{I}}$ and probability measures on each V_i , which govern random walks on it. We consider a transient random walk on the free product arising naturally from the random walks on the V_i . We prove the existence of the rate of escape with respect to the block length, that is, the speed, at which the random walk escapes to infinity, and furthermore we compute formulas for it. For this purpose, we present three different techniques providing three different, equivalent formulas.

2000 *Mathematics subject classification*: Primary 60G50; Secondary 20E06, 60B15.

Keywords and phrases: Random Walks, Free Products, Rate of Escape.

1. Introduction

Consider a transient Markov chain $(Z_n)_{n \in \mathbb{N}_0}$ on a state space V and a suitable length function l on V representing a ‘word length’ with respect to the starting point of the Markov chain. We are interested in whether the sequence of random variables $l(Z_n)/n$ converges almost surely to a constant, and if so, to compute this constant. If the limit exists, it is called the *rate of escape*, or the *drift with respect to l* . In this paper, we study this question for random walks on general free products.

To outline some background material, on the d -dimensional grid \mathbb{Z}^d , where $d \geq 1$, random walks can be described by the sum of n independent and identically distributed random variables, the increments of n steps. By the weak law of large numbers the limit $\lim_{n \rightarrow \infty} |Z_n|/n$, where $|\cdot|$ is the distance on the grid to the starting point of the random walk, exists almost surely. Furthermore, this limit is positive if the increments have non-zero mean vector.

It is well-known that the rate of escape exists also for transitive random walks on finitely generated groups, where the random walks arise from a probability measure on the group elements. This follows from *Kingman’s subadditive ergodic theorem*; see Kingman [9], Derriennic [3] and Guivarc’h [7]. If l is the metric

of the Cayley graph, then the limit $\lim_{n \rightarrow \infty} l(Z_n)/n$ exists almost surely and is positive. There are many detailed results for random walks on groups and wreath products: Mairesse [13] computed an explicit formula in terms of the unique solution of a system of polynomial equations for the rate of escape of random walks on the braid group. Lyons, Pemantle and Peres [11] gave a lower bound for the rate of escape of inward-biased random walks on lamplighter groups. Dyubina [4] proved that the drift on the wreath product $A \wr \mathbb{Z}/2$ is zero, where A is a finitely generated group. Erschler [5] investigated asymptotics of the drift of symmetric random walks on finitely generated groups. An important link between drifts and harmonic analysis was obtained by Varopoulos [19]. He proved that for symmetric finite range random walks on groups the existence of non-trivial bounded harmonic functions is equivalent to a non-zero rate of escape. This leads to a link between the rate of escape and the entropy of random walks, compare e.g. with Kaimanovich and Vershik [8] and Erschler [5]. The rate of escape has also been studied on trees: Cartwright, Kaimanovich and Woess [1] investigated the boundary of homogeneous trees and the drift on them. Nagnibeda and Woess [16, Section 5] proved that the rate of escape of random walks on trees with finitely many cone types is non-zero and give a formula for it.

For a restricted class of free products of finite groups, Mairesse [12] and Mairesse and Mathéus [14] have developed a specific technique for computation of the above limit with respect to the word length. These papers were the starting point for the present investigation of arbitrary free products. We consider the free product of finitely many sets, on which Markov chains are given, and construct in a natural way a random walk on the free product. The techniques we use for rewriting probability generating functions in terms of functions on the factors of the free product were introduced independently and simultaneously by Cartwright and Soardi [2], Woess [22], Voiculescu [20] and McLaughlin [15].

Our aim is to show the existence of the above rate of escape ℓ with respect to the word length, and also to compute formulas for it. For this purpose, we will present three different, equivalent formulas for ℓ using three different techniques. In Section 3 we prove existence and a formula for ℓ by purely probabilistic reasoning. In Section 4 we compute the proposed limit using double generating functions and applying a theorem of Sawyer and Steger [17, Theorem 2.2]. The third approach for the computation of ℓ in Section 5 works only for free products of finitely generated groups and is based on a technique which was already used by Ledrappier [10] and Furstenberg [6]. Section 6 presents sample computations and in Section 7 we give additional remarks about extensions of these techniques to further results.

2. Free Products

2.1. Free Products and Random Walks Let $\mathcal{I} := \{1, \dots, r\}$, $r \geq 2$. Consider r random walks with transition matrices P_i on pairwise disjoint finite or countable state spaces V_i , where $i \in \mathcal{I}$. The corresponding single and n -step transition probabilities are denoted by $p_i(x, y)$ and $p_i^{(n)}(x, y)$, where $x, y \in V_i$. For every $i \in \mathcal{I}$ we select an element o_i of V_i as the ‘root’. To help visualize

this, we think of graphs X_i with vertex sets V_i and roots o_i such that there is an oriented edge $x \rightarrow y$ if and only if $p_i(x, y) > 0$. Furthermore, we shall assume that for every $i \in \mathcal{I}$ and every $x \in V_i$ there is an $n \in \mathbb{N}$ such that $p_i^{(n)}(o_i, x) > 0$. For the sake of simplicity we assume $p_i(x, x) = 0$ for every $i \in \mathcal{I}$ and $x \in V_i$.

Let $V_i^\times := V_i \setminus \{o_i\}$ for every $i \in \mathcal{I}$. The next step is the construction of a new Markov chain on the *free product* $V := V_1 * \dots * V_r$, the set of ‘words’

$$(1) \quad x = x_1 x_2 \dots x_n$$

with letters, also called *blocks*, from the sets V_i^\times such that no two successive letters come from the same V_i . The empty word o describes the root of V . If $u = u_1 \dots u_m \in V$ and $v = v_1 \dots v_n \in V$ with $u_m \in V_i$ and $v_1 \notin V_i$, then uv stands for their concatenation as words. We also define $vo = ov = v$ for all $v \in V$. We regard each V_i as a subset of V , identifying each o_i with o .

We lift P_i to a transition matrix \bar{P}_i on V : if $z \in V$ is o or has last letter not in V_i , and if $v, w \in V_i$, then we set $\bar{p}_i(zv, zw) := p_i(v, w)$. Otherwise we set $\bar{p}_i(x, y) := 0$. We choose $0 < \alpha_1, \dots, \alpha_r \in \mathbb{R}$ with $\sum_{i \in \mathcal{I}} \alpha_i = 1$. Then we obtain a new transition matrix on V given by

$$P = \sum_{i \in \mathcal{I}} \alpha_i \bar{P}_i.$$

The random walk governed by P is described by the sequence of random variables $(Z_n)_{n \in \mathbb{N}_0}$. The associated single and n -step transition probabilities are denoted by $p(x, y)$ and $p^{(n)}(x, y)$ for $x, y \in V$.

Let $x = x_1 \dots x_m \in V \setminus \{o\}$. The **type** $\tau(x)$ of x is defined to be i if $x_m \in V_i$. The **block length** $\ell(x)$ of x is defined to be m . We also set $\tau(o) := 0$ and $\ell(o) := 0$. We want to show existence of the P -almost sure limit $\ell = \lim_{n \rightarrow \infty} \ell(Z_n)/n$, the *rate of escape with respect to the block length*, and to present formulas for it. Let $k \in \mathbb{N}$. Then $x^{(k)} := x_1 \dots x_j$, where $j = \min\{k, m\}$, is the **truncation** at length k . We also write $x^\perp := x_1 \dots x_{m-1}$ for the truncation at length $\ell(x) - 1$, if $x \neq o$. Furthermore we denote by $\tilde{x} := x_m$ the **terminal block** of x , if $m > 0$, and set $\tilde{o} := o$. The **cone** rooted at x is the set

$$C_x := \{y \in V \mid y^{(m)} = x\} \subseteq V.$$

If $y \in V_i$, then the set of **successors** of y is given by

$$\mathcal{S}(y) := \{w \in V_i \mid p_i(y, w) > 0\}$$

and the set of **predecessors** by

$$\mathcal{P}(y) := \{w \in V_i \mid p_i(w, y) > 0\}.$$

We now introduce some probability generating functions. For this purpose, let $T_y := \min\{k \geq 0 \mid Z_k = y\}$, resp. $S_y := \min\{k > 0 \mid Z_k = y\}$, be the stopping time of the first visit, resp. the first return to $y \in V$. Denote by \mathbb{P}_x the probability measure on $V^{\mathbb{N}_0}$ that governs the random walk starting at $x \in V$. For

$z \in \mathbb{C}$ and $x, y \in V$ let

$$\begin{aligned} G(x, y|z) &= \sum_{n \geq 0} p^{(n)}(x, y) z^n, & F(x, y|z) &= \sum_{n \geq 0} \mathbb{P}_x[T_y = n] z^n, \\ U(x, y|z) &= \sum_{n \geq 1} \mathbb{P}_x[S_y = n] z^n, \\ L(x, y|z) &= \sum_{n \geq 0} \mathbb{P}_x[\forall k \in \{1, \dots, n-1\} : Z_k \neq x, Z_n = y] z^n. \end{aligned}$$

The analogous functions for the random walks on the single factors V_i are denoted by $G_i(u, v|z)$, $F_i(u, v|z)$ and $L_i(u, v|z)$, where $u, v \in V_i$. We make the basic assumption that the radius of convergence of $G(o, o|z)$ is greater than 1, which implies transience of our random walk on V . Thus, we may exclude the case $r = 2 = |V_1| = |V_2|$. This convergence property is fulfilled if each p_i is reversible and due to non-amenability also for random walks on free products of finitely generated groups, where the P_i depend only on a probability measure on the single groups. Note that for $|z| < 1$, $\sum_{y \in V_i} G_i(x, y|z) = 1/(1-z)$ for every $i \in \mathcal{I}$ and all $x \in V_i$. This will be used several times in the sequel.

Recall the following equations:

LEMMA 2.1. *Let $x, y \in V$, $w \in V \setminus \{o\}$ such that $w^{(1)} \notin V_{\tau(x)}$, and $z \in \mathbb{C}$. Then:*

$$\begin{aligned} (i) \quad G(x, x|z) &= \frac{1}{1 - U(x, x|z)}, \\ (ii) \quad G(x, y|z) &= F(x, y|z) \cdot G(y, y|z), \\ (iii) \quad G(x, y|z) &= G(x, x|z) \cdot L(x, y|z), \\ (iv) \quad F(o, xw|z) &= F(o, x|z) \cdot F(x, xw|z), \\ (v) \quad L(o, xw|z) &= L(o, x|z) \cdot L(x, xw|z), \\ (vi) \quad L(x, xw|z) &= L(o, w|z). \end{aligned}$$

Equations analogous to (i), (ii), (iii) hold for the generating functions on the single factors V_i for every $i \in \mathcal{I}$.

PROOF. For (i) and (ii) see Woess [23, Lemma 1.13]. Equation (iii) is obtained by conditioning with respect to the last visit at x before finally walking to y . (iv) and (v) are obtained by conditioning with respect to the first resp. last visit at x , which must be visited before finally walking to xw . (vi) holds, as due to the tree-like structure of the free product the probability of walking from x to xw in n steps without returning to x is the same as walking from o to w in n steps without returning to o . \square

We now explain the correspondence between $F(x, y|z)$ and $F_i(x, y|z)$, resp. $L(x, y|z)$ and $L_i(x, y|z)$. Therefore define for $i \in \mathcal{I}$ and $z \in \mathbb{C}$

$$\bar{H}_i(z) := \sum_{n=2}^{\infty} \mathbb{P}_o[S_o = n, Z_1 \notin V_i] z^n \quad \text{and} \quad \xi_i(z) := \frac{\alpha_i z}{1 - \bar{H}_i(z)};$$

see Woess [23, Proposition 9.18]. Note that $\bar{H}_i(1)$ is the probability of starting at some $x \in V_i$ and returning to the same x without having visited a neighbour

$y \in \mathcal{S}(x)$ of x . Similarly $\xi_i(1)$ is the probability of starting at some $x \in V_i$ and at some time visiting a neighbour $y \in \mathcal{S}(x)$. Observe that for positive z the functions $\bar{H}_i(z)$ and $\xi_i(z)$ are strictly increasing inside their radii of convergence, which are greater than 1.

LEMMA 2.2. *Let $i \in \mathcal{I}$, $x, y \in V_i$ and $z \in \mathbb{C}$. Then*

$$(i) \ F(x, y|z) = F_i(x, y|\xi_i(z)) \quad \text{and} \quad (ii) \ L(x, y|z) = L_i(x, y|\xi_i(z)).$$

For the proof of (i) see Woess [23, Proposition 9.18 (c)]. Statement (ii) is proved analogously.

LEMMA 2.3. $\xi_i := \xi_i(1) < 1$ for all $i \in \mathcal{I}$.

PROOF. Let $H_i(z) := U(o, o|z) - \bar{H}_i(z)$. By transience we have

$$U(o, o|1) = \sum_{i \in \mathcal{I}} H_i(1) < 1.$$

Furthermore

$$H_i(1) = \alpha_i \sum_{s \in \mathcal{S}(o_i)} p_i(o_i, s) \underbrace{F(s, o|1)}_{\leq 1} \leq \alpha_i.$$

Hence,

$$\xi_i = \frac{\alpha_i}{1 - \sum_{j \in \mathcal{I} \setminus \{i\}} H_j(1)} \leq \frac{\alpha_i}{1 - \sum_{j \in \mathcal{I} \setminus \{i\}} \alpha_j} = \frac{\alpha_i}{1 - (1 - \alpha_i)} = 1.$$

Observe that $\xi_j < 1$ for all $j \in \mathcal{I} \setminus \{i\}$, if $H_i(1) < \alpha_i$ for some $i \in \mathcal{I}$. Assume $H_i(1) = \alpha_i$ for some $i \in \mathcal{I}$. Then $\bar{H}_i(1) = U(o, o|1) - H_i(1) < 1 - \alpha_i$, and thus there is $j \in \mathcal{I} \setminus \{i\}$ such that $H_j(1) < \alpha_j$. Thus $\xi_i < 1$. Since

$$H_i(1) = \alpha_i \sum_{s \in \mathcal{S}(o_i)} p_i(o_i, s) F(s, o|1) = \alpha_i$$

we have $F(s, o|1) = 1$ for all $s \in \mathcal{S}(o_i)$. But now we obtain the contradiction

$$F(s, o|1) = F_i(s, o_i|\xi_i) < 1,$$

as $\xi_i < 1$ and $F_i(s, o_i|x)$, $0 \leq x \in \mathbb{R}$, is strictly increasing with $F_i(s, o_i|1) \leq 1$. \square

2.2. Limit of the Random Walk As we have assumed transience for the random walk on V , the random walk escapes to infinity in the sense that almost surely every finite $A \subseteq V$ is visited only finitely often. We shall now investigate the route of the escape of the random walk on V , which provides the main tool for further computations. Define for $x \in V$, $i \in \mathcal{I}$ with $\tau(x) \neq i$ and $S \subseteq V_i$ the set $xS := \{xy \mid y \in S\}$. Then we obtain:

LEMMA 2.4. *If $x \in V$ and $\tau(x) \neq i$ then*

$$\mathbb{P}_o[Z_n \in xV_i \text{ holds for infinitely many } n] = 0.$$

PROOF. By Lemmas 2.1 and 2.2,

$$\begin{aligned}
\sum_{n \geq 0} \mathbb{P}_o[Z_n \in xV_i] &= \sum_{y \in V_i} G(o, xy|1) \\
&= \sum_{y \in V_i} G(o, o|1) L(o, x|1) L_i(o_i, y|\xi_i) \\
&= G(o, o|1) \cdot L(o, x|1) \cdot \sum_{y \in V_i} \frac{G_i(o_i, y|\xi_i)}{G_i(o_i, o_i|\xi_i)} \\
&= \frac{G(o, o|1) L(o, x|1)}{G_i(o_i, o_i|\xi_i)} \frac{1}{1 - \xi_i} < \infty.
\end{aligned}$$

The Borel-Cantelli lemma implies the proposed statement. \square

Now we are able to specify how the random walk on V escapes to infinity. Let V_∞ denote the set of infinite words $x_1x_2 \dots$ in which each of the letters x_j belongs to $\bigcup_{i \in \mathcal{I}} V_i^\times$, no consecutive letters come from the same V_i^\times , and infinitely many letters come from each V_i^\times . Then we obtain:

PROPOSITION 2.5. $\ell(Z_n)$ tends to infinity \mathbb{P}_o -a.s., as $n \rightarrow \infty$. Furthermore, there exists a V_∞ -valued random variable Z_∞ , such that

$$\lim_{n \rightarrow \infty} Z_n = Z_\infty \quad \mathbb{P}_o - a.s.,$$

with convergence in the sense that the length of the common prefix of Z_n and Z_∞ tends to infinity.

PROOF. We prove by induction that for each $m \in \mathbb{N}$ there is almost surely some $n_m \in \mathbb{N}$ with $\ell(Z_{n_m}) = m$ and $\ell(Z_n) > m$ for all $n > n_m$. By Lemma 2.4, the random walk visits the state set $\bigcup_{i \in \mathcal{I}} V_i$ finitely often \mathbb{P}_o -a.s.. Therefore there is almost surely some $n_1 \in \mathbb{N}$ such that $Z_{n_1} \in \bigcup_{i \in \mathcal{I}} V_i$ \mathbb{P}_o -a.s. and $Z_n \notin \bigcup_{i \in \mathcal{I}} V_i$ for all $n > n_1$. Thus $\ell(Z_{n_1}) = 1$ and $\ell(Z_n) > 1$ for all $n > n_1$.

Assume now that $\ell(Z_{n_m}) = m$ and $\ell(Z_n) > m$ for all $n > n_m$ and let $\mathcal{I}' := \mathcal{I} \setminus \{\tau(Z_{n_m})\}$. Again by Lemma 2.4 the random walk visits the state set $\bigcup_{i \in \mathcal{I}'} Z_{n_m} V_i$ finitely often \mathbb{P}_o -a.s.. Then there is almost surely some $n_{m+1} \in \mathbb{N}$ such that $Z_{n_{m+1}} \in \bigcup_{i \in \mathcal{I}'} Z_{n_m} V_i$ and $Z_n \notin \bigcup_{i \in \mathcal{I}'} Z_{n_m} V_i$ for all $n > n_{m+1}$. Thus $\ell(Z_{n_{m+1}}) = m + 1$ and $\ell(Z_n) > m + 1$ for all $n > n_{m+1}$. Thus $\ell(Z_n)$ tends to infinity, as $n \rightarrow \infty$.

Obviously the sequence $(Z_n)_{n \in \mathbb{N}_0}$ converges to an infinite word in V_∞ with $Z_\infty^{(m)} = Z_{n_m}$ for all $m \in \mathbb{N}$. \square

3. Exit Time Technique

In this section we investigate the random walk on V in detail, prove the existence of ℓ and derive a formula for it. The following technique was motivated by

Nagnibeda and Woess [16, Section 5]. Let $k \in \mathbb{N}$. The exit time with respect to the block length k is

$$\mathbf{e}_k := \min \left\{ m \in \mathbb{N} \mid \forall n \geq m : Z_n^{(k)} \text{ constant} \right\}.$$

In particular $\mathbf{e}_0 = 0$. The exit point with respect to the block length k is $W_k := Z_{\mathbf{e}_k}$. Thus, \mathbf{e}_k is the first instant from which point the first k blocks remain constant, and $W_k = x$ if and only if at time $\mathbf{e}_k - 1$ the random walk is at state $x_1 \dots x_{k-1}$ with some $s \in \mathcal{P}(x_k)$, at time \mathbf{e}_k at state x , and thereafter remains in the cone C_x .

As Z_n converges almost surely to a random variable Z_∞ with values in V_∞ , we have $\mathbf{e}_k \rightarrow \infty$ as $k \rightarrow \infty$, almost surely. The k -th increment is $\mathbf{i}_k := \mathbf{e}_k - \mathbf{e}_{k-1}$ and the maximal temporary exit time at time n is defined as

$$\mathbf{k}(n) := \max \{ k \in \mathbb{N}_0 \mid \mathbf{e}_k \leq n \}.$$

Define now for $i \in \mathcal{I}$, $y \in V$ and $n \in \mathbb{N}_0$

$$k_i^{(n)}(o, y) := \mathbb{P}_o[\forall j \in \{0, \dots, n\} : Z_j \notin V_i^\times, Z_n = y]$$

and the corresponding generating function

$$(2) \quad K_i(o, y|z) := \sum_{n=0}^{\infty} k_i^{(n)}(o, y) z^n = \sum_{n \geq 0} \bar{H}_i(z)^n \cdot L(o, y|z) = \frac{L(o, y|z)}{1 - \bar{H}_i(z)}.$$

Now we can prove:

PROPOSITION 3.1. $(W_k, \mathbf{i}_k)_{k \in \mathbb{N}}$ is a Markov chain with transition probabilities

$$\begin{aligned} & \mathbb{P}_o[W_{k+1} = w_{k+1}, \mathbf{i}_{k+1} = n_{k+1} \mid W_k = w_k, \mathbf{i}_k = n_k] \\ &= \frac{1 - \xi_{\tau(w_{k+1})}}{1 - \xi_{\tau(w_k)}} \cdot \sum_{s \in \mathcal{P}(y)} \left[k_{\tau(w_k)}^{(n_{k+1}-1)}(o, s) \cdot p(s, y) \right] \end{aligned}$$

for $n_k, n_{k+1} \in \mathbb{N}$, $w_k = x_1 \dots x_k \in V$, $w_{k+1} = w_k y$, where $y \in \bigcup_{i \in \mathcal{I} \setminus \{\tau(w_k)\}} V_i^\times$.

PROOF. Define $\bar{V}_i = \bigcup_{j \in \mathcal{I} \setminus \{i\}} V_j^\times$. Let $w_0 = o$, $w_1 = g_1 \in \bigcup_{i \in \mathcal{I}} V_i^\times$ and $w_i = w_{i-1} g_i$ with $g_i \in \bar{V}_{\tau(w_{i-1})}$ for $2 \leq i \leq k$.

For $i \in \{1, \dots, k\}$ the inclusion $[W_{i+1} = w_{i+1}] \subseteq [W_i = w_i]$ holds, as w_{i+1} determines the element w_i uniquely. Let $n_1, \dots, n_{k+1} \in \mathbb{N}$ and write for $m \in \{k, k+1\}$

$$[W_1^m = w_1^m, \mathbf{i}_1^m = n_1^m] := [\forall j \in \{1, \dots, m\} : W_j = w_j, \mathbf{i}_j = n_j].$$

This event can be described as follows: start at o , walk in $n_1 - 1$ steps to a predecessor of w_1 inside $V_{\tau(w_1)}$, then walk to w_1 ; then stay inside C_{w_1} and walk in $n_2 - 1$ steps to a vertex in $w_1 \mathcal{P}(g_2)$, from there to w_2 , and so on. More formally we obtain, writing $n_1^s = \sum_{t=1}^s n_t$:

$$\begin{aligned} & \mathbb{P}_o[W_1^k = w_1^k, \mathbf{i}_1^k = n_1^k] \\ &= \mathbb{P}_o \left[\begin{array}{c} \forall \lambda \in \{0, \dots, k-1\} \forall j \in \{1, \dots, n_\lambda - 1\} : \\ Z_{n_1^\lambda + j} \in C_{w_\lambda}, Z_{n_1^{\lambda+1} - 1} \in w_\lambda \mathcal{P}(g_{\lambda+1}), Z_{n_1^{\lambda+1}} = w_{\lambda+1} \end{array} \right] \cdot (1 - \xi_{\tau(w_k)}). \end{aligned}$$

Analogously,

$$\begin{aligned} & \mathbb{P}_o[W_1^{k+1} = w_1^{k+1}, \mathbf{i}_1^{k+1} = n_1^{k+1}] \\ &= \mathbb{P}_o \left[\begin{array}{c} \forall \lambda \in \{0, \dots, k-1\} \forall j \in \{1, \dots, n_\lambda - 1\} : \\ Z_{n_1^\lambda + j} \in C_{w_\lambda}, Z_{n_1^{\lambda+1} - 1} \in w_\lambda \mathcal{P}(g_{\lambda+1}), Z_{n_1^{\lambda+1}} = w_{\lambda+1} \end{array} \right] \\ & \quad \mathbb{P}_{w_k} \left[\begin{array}{c} \forall j \in \{1, \dots, n_{k+1} - 2\} : Z_j \in C_{w_k}, \\ Z_{n_{k+1} - 1} \in w_k \mathcal{P}(g_{k+1}), Z_{n_{k+1}} = w_{k+1} \end{array} \right] \cdot (1 - \xi_{\tau(w_{k+1})}). \end{aligned}$$

Thus we obtain the conditional probabilities:

$$\begin{aligned} & \mathbb{P}_o[W_{k+1} = w_{k+1}, \mathbf{i}_{k+1} = n_{k+1} \mid W_1^k = w_1^k, \mathbf{i}_1^k = n_1^k] \\ &= \frac{1 - \xi_{\tau(w_{k+1})}}{1 - \xi_{\tau(w_k)}} \cdot \mathbb{P}_{w_k} \left[\begin{array}{c} \forall j \in \{1, \dots, n_{k+1} - 2\} : Z_j \in C_{w_k}, \\ Z_{n_{k+1} - 1} \in w_k \mathcal{P}(g_{k+1}), Z_{n_{k+1}} = w_{k+1} \end{array} \right] \\ &= \frac{1 - \xi_{\tau(w_{k+1})}}{1 - \xi_{\tau(w_k)}} \cdot \sum_{s \in \mathcal{P}(y)} \left[k_{\tau(w_k)}^{(n_{k+1}-1)}(o, s) \cdot p(s, y) \right]. \end{aligned}$$

□

Observe that the transition probabilities of the stochastic process $(W_k, \mathbf{i}_k)_{k \in \mathbb{N}}$ depend only on $\tau_k := \tau(W_k)$, $\tau(W_{k+1})$ and \widetilde{W}_{k+1} . Hence, the stochastic process $(\widetilde{W}_k, \mathbf{i}_k, \tau_k)_{k \in \mathbb{N}}$ is an irreducible Markov chain on the state space

$$\mathcal{A} := \left\{ (y, n, j) \mid j \in \mathcal{I}, y \in V_j^\times, n \in \mathbb{N}, \exists i \in \mathcal{I} \setminus \{j\} \exists s \in \mathcal{P}(y) : k_i^{(n-1)}(o, s) > 0 \right\}$$

with transition probabilities

$$q((x, m, i), (y, n, j)) = \begin{cases} 0, & \text{if } i = j \\ \frac{1 - \xi_j}{1 - \xi_i} \cdot \sum_{s \in \mathcal{P}(y)} \left[k_i^{(n-1)}(o, s) \cdot p(s, y) \right], & \text{if } i \neq j \end{cases}.$$

For convenience, we write $q((x, m, i), (y, n, j)) := 0$, if $j \in \mathcal{I}$, $y \in V_j$, but $(y, n, j) \notin \mathcal{A}$. As the probabilities $q((x, m, i), (y, n, j))$ do not depend on x and m , the sequence $(\tau_k)_{k \in \mathbb{N}}$ is also a Markov chain on the state space \mathcal{I} with transition probabilities

$$(3) \quad \hat{q}(i, j) := \sum_{y \in V_j^\times} \sum_{n \geq 1} q((x, m, i), (y, n, j))$$

for $i, j \in \mathcal{I}$ with $i \neq j$ and $\hat{q}(i, i) = 0$. Note that $x \in V_i$ and $m \in \mathbb{N}$ can be chosen arbitrarily, such that $(x, m, i) \in \mathcal{A}$. As \mathcal{I} is finite, $(\tau_k)_{k \in \mathbb{N}}$ possesses an invariant probability measure $\nu : \mathcal{I} \rightarrow [0, 1]$, that is, for every $j \in \mathcal{I}$

$$(4) \quad \sum_{i \in \mathcal{I}} \nu(i) \cdot \hat{q}(i, j) = \nu(j).$$

We now define for $j \in \mathcal{I}$, $y \in V_j^\times$ and $n \in \mathbb{N}$

$$\pi(y, n, j) := \sum_{i \in \mathcal{I}} \nu(i) \cdot q((x, m, i), (y, n, j)),$$

which is an invariant probability measure of the stochastic process $(\widetilde{W}_k, \mathbf{i}_k, \tau_k)_{k \in \mathbb{N}}$, that is,

$$\sum_{(x,m,i) \in \mathcal{A}} \pi(x, m, i) q((x, m, i), (y, n, j)) = \pi(y, n, j)$$

holds for all $(y, n, j) \in \mathcal{A}$.

PROPOSITION 3.2. *There is a number $\Lambda \in \mathbb{R}$, such that*

$$\frac{\mathbf{e}_k}{k} \xrightarrow{k \rightarrow \infty} \Lambda \quad \mathbb{P}_o - a.s..$$

PROOF. Consider the function $g : \mathcal{A} \rightarrow \mathbb{N}$, $(y, n, j) \mapsto n$. An application of the ergodic theorem for positive recurrent Markov chains shows that

$$\frac{1}{k} \sum_{l=1}^k g(\widetilde{W}_l, \mathbf{i}_l, \tau_l) = \frac{\mathbf{e}_k - \mathbf{e}_0}{k} = \frac{\mathbf{e}_k}{k} \xrightarrow{k \rightarrow \infty} \int g d\pi \quad \mathbb{P}_o - a.s.,$$

if $\int g d\pi < \infty$ holds. Hence it is sufficient to show finiteness of this integral. Noting that

$$\begin{aligned} \int g d\pi &= \sum_{i \in \mathcal{I}} \nu(i) \cdot \sum_{(y,n,j) \in \mathcal{A}} n \cdot q((x, m, i), (y, n, j)) \\ &= \sum_{i \in \mathcal{I}} \frac{\nu(i)}{1 - \xi_i} \sum_{j \in \mathcal{I} \setminus \{i\}} \alpha_j (1 - \xi_j) \underbrace{\sum_{n \geq 1} n \sum_{y \in V_j^\times} \sum_{s \in \mathcal{P}(y)} k_i^{(n-1)}(o, s) \cdot p_j(s, y)}_{(*)}, \end{aligned}$$

we now interpret the sum $(*)$ as a power series evaluated at 1. We have

$$\begin{aligned} &\sum_{n \geq 1} n \sum_{y \in V_j^\times} \sum_{s \in \mathcal{P}(y)} k_i^{(n-1)}(o, s) p_j(s, y) z^{n-1} \\ &= \frac{\partial}{\partial z} \left[\underbrace{\sum_{n \geq 1} \sum_{y \in V_j^\times} \sum_{s \in \mathcal{P}(y)} k_i^{(n-1)}(o, s) p_j(s, y) z^n}_{=: \gamma_{i,j}(z)} \right]. \end{aligned}$$

Now it is sufficient to show that the sum $\gamma_{i,j}(z)$ has radius of convergence $R_{i,j} > 1$ for all $i, j \in \mathcal{I}$ with $i \neq j$:

$$\begin{aligned} \gamma_{i,j}(z) &= \sum_{n \geq 1} \sum_{y \in V_j^\times} \sum_{s \in \mathcal{P}(y)} k_i^{(n-1)}(o, s) \cdot p_j(s, y) \cdot z^n \\ &= \sum_{n \geq 1} \sum_{s \in V_j} k_i^{(n-1)}(o, s) \cdot z^n - \sum_{n \geq 1} \sum_{s \in \mathcal{P}(o_j)} k_i^{(n-1)}(o, s) \cdot p_j(s, o_j) \cdot z^n \\ &= z \cdot \underbrace{\sum_{y \in V_j} K_i(o, y|z)}_{(**)} - z \cdot \underbrace{\sum_{s \in \mathcal{P}(o_j)} K_i(o, s|z) \cdot p_j(s, o_j)}_{(***)}. \end{aligned}$$

From Equation (2) we obtain

$$\sum_{y \in V_j} K_i(o, y|z) = \sum_{y \in V_j} \frac{L(o, y|z)}{1 - \bar{H}_i(z)} = \frac{1}{1 - \bar{H}_i(z)} \sum_{y \in V_j} L_j(o_j, y|\xi_j(z)),$$

and also

$$\begin{aligned} \sum_{y \in V_j} L_j(o_j, y|\xi_j(z)) &= \frac{1}{G_j(o_j, o_j|\xi_j(z))} \sum_{y \in V_j} G_j(o_j, y|\xi_j(z)) \\ &= \frac{1}{G_j(o_j, o_j|\xi_j(z))} \cdot \frac{1}{1 - \xi_j(z)}. \end{aligned}$$

Thus the sum (**) has radius of convergence greater than 1. Furthermore, by (2) and Lemmas 2.1 and 2.2,

$$\begin{aligned} &\sum_{s \in \mathcal{P}(o_j)} p_j(s, o_j) K_i(o, s|z) \\ &= \frac{1}{1 - \bar{H}_i(z)} \sum_{s \in \mathcal{S}(o_j)} p_j(s, o_j) \frac{G_j(o_j, s|\xi_j(z))}{G_j(o_j, o_j|\xi_j(z))} \\ &= \frac{1}{(1 - \bar{H}_i(z)) \cdot G_j(o_j, o_j|\xi_j(z))} \sum_{s \in \mathcal{P}(o_j)} p_j(s, o_j) \xi_j(z) G_j(o_j, s|\xi_j(z)) \frac{1}{\xi_j(z)} \\ &= \frac{G_j(o_j, o_j|\xi_j(z)) - 1}{(1 - \bar{H}_i(z)) \cdot G_j(o_j, o_j|\xi_j(z)) \cdot \xi_j(z)}. \end{aligned}$$

Thus the sum (***) has radius of convergence greater than 1, and also $\gamma_{i,j}(z)$, whence $\int g d\pi$ is finite. \square

Using the above, we can rewrite $\gamma_{i,j}(z)$ as

$$(5) \quad \gamma_{i,j}(z) = \frac{1}{\alpha_i} \cdot \frac{\xi_i(z)}{\xi_j(z)} \cdot \left(\frac{1}{(1 - \xi_j(z)) \cdot G_j(o_j, o_j|\xi_j(z))} - 1 \right)$$

and therefore

$$(6) \quad \Lambda = \sum_{\substack{i, j \in \mathcal{I}, \\ i \neq j}} \nu(i) \cdot \alpha_j \cdot \frac{1 - \xi_j}{1 - \xi_i} \cdot \gamma'_{i,j}(1).$$

The following theorem is now obtained precisely as in Nagnibeda and Woess [16, proof of Theorem D]:

THEOREM 3.3.

$$\frac{\ell(Z_n)}{n} \xrightarrow{n \rightarrow \infty} \ell = \frac{1}{\Lambda} \quad \mathbb{P}_o - a.s.$$

Finally, we show how to compute the invariant probability measure ν explicitly. For this purpose, it is sufficient to compute the transition probabilities $\hat{q}(i, j)$ for all $i, j \in \mathcal{I}$. By solving the system of linear equations given by Equations (4) ν is obtained. The next lemma proposes a formula for $\hat{q}(i, j)$:

LEMMA 3.4. *Let $i, j \in \mathcal{I}$ with $i \neq j$. Then $\hat{q}(i, i) = 0$ and*

$$\hat{q}(i, j) = \frac{\alpha_j}{\alpha_i} \cdot \frac{\xi_i}{\xi_j} \cdot \frac{1 - \xi_j}{1 - \xi_i} \cdot \left(\frac{1}{(1 - \xi_j) G_j(\xi_j)} - 1 \right), \quad \text{where } G_j(\xi_j) := G_j(o_j, o_j | \xi_j).$$

PROOF. By definition of W_k and $\mathbb{P}_o[Z_\infty \in V_\infty] = 1$ it follows that $\hat{q}(i, i) = 0$. Considering (3) and the computations in the proof of Proposition 3.2 leads to $\hat{q}(i, j) = (1 - \xi_j)/(1 - \xi_i) \cdot \alpha_j \cdot \gamma_{i,j}(1)$. \square

We now give an explicit formula for ν :

$$(7) \quad \nu(i) = c \cdot \frac{\alpha_i (1 - \xi_i)}{\xi_i} \cdot (1 - (1 - \xi_i) G_i(\xi_i)),$$

where $c > 0$ is chosen so that $\sum_{i \in \mathcal{I}} \nu(i) = 1$. This is indeed an invariant measure, because, writing $x(i) = 1 - (1 - \xi_i) G_i(\xi_i)$, the invariance condition on ν is just

$$\sum_{i \in \mathcal{I} \setminus \{j\}} x(i) = \frac{x(j)}{\frac{1}{(1 - \xi_j) G_j(\xi_j)} - 1} \quad \text{for each } j \in \mathcal{I}$$

or, equivalently, that

$$\sum_{i \in \mathcal{I}} x(i) = 1.$$

The following lemma verifies that this equation holds.

LEMMA 3.5. *Let $i \in \mathcal{I}$. Then*

$$\rho(i) := \mathbb{P}_o \left[Z_\infty^{(1)} \in V_i^\times, \forall n \in \mathbb{N} : Z_n \notin \bigcup_{j \in \mathcal{I} \setminus \{i\}} V_j \right] = \frac{1 - (1 - \xi_i) G_i(\xi_i)}{G(o, o | 1)}.$$

PROOF. By transience, o is visited only finitely often \mathbb{P}_e -a.s., that is,

$$\sum_{i \in \mathcal{I}} G(o, o | 1) \rho(i) = 1.$$

This yields

$$\begin{aligned} \rho(i) &= \sum_{y \in V_i^\times} \sum_{j \in \mathcal{I} \setminus \{i\}} L(o, y | 1) \cdot \rho(j) \\ &= \sum_{y \in V_i^\times} L(o, y | 1) \cdot (G(o, o | 1)^{-1} - \rho(i)) \\ &= (G(o, o | 1)^{-1} - \rho(i)) \sum_{y \in V_i^\times} \frac{G_i(o_i, y | \xi_i)}{G_i(\xi_i)} \\ &= (G(o, o | 1)^{-1} - \rho(i)) \cdot \left(\frac{1}{(1 - \xi_i) \cdot G_i(\xi_i)} - 1 \right). \end{aligned}$$

This leads to the proposed equation. \square

We summarize this section: if we know for each factor V_i , $i \in \mathcal{I}$, the first visit generating function $F_i(x, o_i|z)$, when starting at a predecessor $x \in \mathcal{P}(o_i)$ of o_i , and the Green function $G_i(o_i, o_i|z)$, it is possible to compute $\xi_i(z)$ by solving a finite system of characteristic equations. This is in fact only possible when the generating functions are known and not too complicated. The measure ν can then be computed by (7) and then ℓ found using $\ell = 1/\Lambda$ and (6). Sample computations are presented in Section 6.

4. Double Generating Functions

In this section we compute the rate of escape ℓ for the random walk on V using double generating functions. The main tool for our computation is the following theorem:

THEOREM 4.1 (Sawyer and Steger). *Let (Y_n) be a sequence of real-valued random variables such that for some $\delta > 0$,*

$$\mathbb{E}\left(\sum_{n \geq 0} \exp(-rY_n - sn)\right) = \frac{C(r, s)}{g(r, s)} \quad \text{for } 0 < r, s < \delta,$$

where $C(r, s)$ and $g(r, s)$ are analytic for $|r|, |s| < \delta$ and $C(0, 0) \neq 0$. Denote by g_r and g_s the partial derivatives of g with respect to r and s . Then

$$\frac{Y_n}{n} \xrightarrow{n \rightarrow \infty} \ell = \frac{g_r(0, 0)}{g_s(0, 0)} \quad \text{almost surely.}$$

For the proof, see Sawyer and Steger [17, Theorem 2.2].

Setting $Y_n = \ell(Z_n)$, $w = e^{-r}$ and $z = e^{-s}$, to find ℓ in our context it is sufficient to investigate the double generating function

$$\mathcal{E}(w, z) := \sum_{x \in V} \sum_{n \geq 0} p^{(n)}(o, x) w^{\ell(x)} z^n = \sum_{x \in V} G(o, x|z) w^{\ell(x)}$$

and to apply Theorem 4.1. To this end, introduce further double generating functions. Write $V^\times := V \setminus \{o\}$ and define

$$\mathcal{L}(w, z) := 1 + \sum_{n \geq 1} \sum_{x=x_1 \dots x_n \in V^\times} \prod_{j=1}^n w L_{\tau(x_j)}(o_{\tau(x_j)}, x_j | \xi_{\tau(x_j)}(z))$$

and for $i \in \mathcal{I}$

$$\mathcal{L}_i^+(w, z) := \sum_{x \in V_i^\times} L_i(o_i, x | \xi_i(z)) w,$$

$$\mathcal{L}_i(w, z) := \mathcal{L}_i^+(w, z) \left(1 + \sum_{n \geq 2} \sum_{\substack{x_2, \dots, x_n \in V^\times \\ \tau(x_2) \neq i}} \prod_{j=2}^n w L_{\tau(x_j)}(o_{\tau(x_j)}, x_j | \xi_{\tau(x_j)}(z)) \right).$$

Thus we have the equation

$$\mathcal{L}(w, z) = 1 + \sum_{i \in \mathcal{I}} \mathcal{L}_i(w, z).$$

If $0 \leq w, z < 1$, the convergence of the series $\mathcal{L}(w, z)$ follows by $\mathcal{L}(w, z) \leq \mathcal{E}(w, z)$. Hence, $\mathcal{L}_i^+(w, z)$ and also $\mathcal{L}_i(w, z)$, $i \in \mathcal{I}$, converge if $|w|, |z| < 1$. The next lemma provides another representation of $\mathcal{L}(w, z)$.

LEMMA 4.2. *Let $w, z \in \mathbb{R}$ with $0 \leq w, z < 1$. Then:*

$$\mathcal{L}(w, z) = \frac{1}{1 - \mathcal{L}^*(w, z)}, \quad \text{where} \quad \mathcal{L}^*(w, z) = \sum_{i \in \mathcal{I}} \frac{\mathcal{L}_i^+(w, z)}{1 + \mathcal{L}_i^+(w, z)}.$$

PROOF. Let $w, z \in \mathbb{R}$ with $0 < w, z < 1$. First we have

$$\mathcal{L}_i(w, z) = \mathcal{L}_i^+(w, z) \cdot \left(1 + \sum_{j \in \mathcal{I} \setminus \{i\}} \mathcal{L}_j(w, z)\right) \quad \text{for all } i \in \mathcal{I},$$

and from convergence of $\mathcal{L}(w, z)$ we get

$$\mathcal{L}_i(w, z) = \mathcal{L}_i^+(w, z) \cdot (\mathcal{L}(w, z) - \mathcal{L}_i(w, z)).$$

As $\mathcal{L}_i^+(w, z) \geq 0$ holds, the last equation is equivalent to

$$\mathcal{L}_i(w, z) = \frac{\mathcal{L}_i^+(w, z)}{1 + \mathcal{L}_i^+(w, z)} \mathcal{L}(w, z).$$

Thus

$$\mathcal{L}(w, z) = 1 + \sum_{i \in \mathcal{I}} \mathcal{L}_i(w, z) = 1 + \sum_{i \in \mathcal{I}} \frac{\mathcal{L}_i^+(w, z)}{1 + \mathcal{L}_i^+(w, z)} \mathcal{L}(w, z).$$

As $\mathcal{L}(w, z) < \infty$, we get

$$\sum_{i \in \mathcal{I}} \frac{\mathcal{L}_i^+(w, z)}{1 + \mathcal{L}_i^+(w, z)} < 1.$$

and the result follows. \square

COROLLARY 4.3. *Let $w, z \in \mathbb{R}$ with $0 \leq w, z < 1$. Then*

$$\mathcal{E}(w, z) = \frac{G(o, o|z)}{1 - \mathcal{L}^*(w, z)}.$$

PROOF. Let $w, z \in \mathbb{R}$ with $0 < w, z < 1$. Applying Lemma 4.2 yields the proposed equation:

$$\begin{aligned} \mathcal{E}(w, z) &= \sum_{x \in V} G(o, o|z) L(o, x|z) w^{\ell(x)} \\ &= G(o, o|z) \left(1 + \sum_{n \geq 1} \sum_{x = x_1 \dots x_n \in V^\times} L(o, x|z) w^{\ell(x)}\right) \\ &= G(o, o|z) \cdot \mathcal{L}(w, z). \end{aligned}$$

\square

We can now conclude and compute a formula for the rate of escape ℓ . Rewriting $\mathcal{L}^*(w, z)$ after some manipulations involving Lemma 2.1 yields

$$\mathcal{L}^*(w, z) = \sum_{i \in \mathcal{I}} \frac{w \left(\frac{1}{1 - \xi_i(z)} - G_i(o_i, o_i | \xi_i(z)) \right)}{\frac{w}{1 - \xi_i(z)} + (1 - w) G_i(o_i, o_i | \xi_i(z))}.$$

Now we define

$$C(w, z) := G(o, o | z) \quad \text{and} \\ g(w, z) := 1 - \sum_{i \in \mathcal{I}} \frac{w \left(\frac{1}{1 - \xi_i(z)} - G_i(o_i, o_i | \xi_i(z)) \right)}{\frac{w}{1 - \xi_i(z)} + (1 - w) G_i(o_i, o_i | \xi_i(z))}.$$

and we have

$$\mathcal{E}(w, z) = \frac{C(w, z)}{g(w, z)} \quad \text{for } 0 \leq w, z < 1.$$

The constraints required for an application of Theorem 4.1 are obviously fulfilled, as $G(o, o | z)$ has radius of convergence greater than 1 and $\xi_i < 1$. We apply it now, where g_w and g_z denote the partial derivatives of g with respect to w and z , respectively. Hence, we can conclude

$$\frac{\ell(Z_n)}{n} \xrightarrow{n \rightarrow \infty} \ell = \frac{g_w(1, 1)}{g_z(1, 1)} \quad \mathbb{P}_o - a.s..$$

Simplifications yield the following formula for ℓ , where we write $\xi_i = \xi_i(1)$ and $G_i(\xi_i) := G_i(o_i, o_i | \xi_i)$:

$$(8) \quad \ell = \frac{\sum_{i \in \mathcal{I}} \left[(1 - (1 - \xi_i) G_i(\xi_i)) \cdot G_i(\xi_i) \cdot (1 - \xi_i) \right]}{\sum_{i \in \mathcal{I}} \left[\xi_i'(1) \cdot (G_i(\xi_i) - (1 - \xi_i) G_i'(\xi_i)) \right]}.$$

Observe that $\xi_i, G_i(\xi_i) > 0$, $\xi_i < 1$ and $G_i(\xi_i) < (1 - \xi_i)^{-1}$. Thus $\ell > 0$.

5. Free Products of Groups

In this section we present a third technique for the computation of the rate of escape ℓ of the block length for the random walk on the free product. This technique is restricted to the case of a free product of *groups*. Therefore let Γ_i , $i \in \mathcal{I}$, be non-trivial finitely generated groups. We assume that the groups have pairwise trivial intersections, but they may be isomorphic. Denote by e_i the identity on Γ_i . The elements of the free product $\Gamma := \Gamma_1 * \dots * \Gamma_r$ are represented as words in the sense of (1) and e is identified with the empty word.

We can define a group operation on Γ : the product of $u, v \in \Gamma$ is the concatenation of the words u and v with possible cancellations and contractions in the middle to get the representative form of the product word. We exclude the case $r = 2 = |\Gamma_1| = |\Gamma_2|$. This ensures that the free group product is non-amenable, yielding that each of our constructed random walks on Γ is transient and $G(e, e | z)$ has radius of convergence greater than 1. (See Woess [23, Theorem

10.10, Proposition 12.4, Corollary 12.5]). We write $\Gamma_i^\times = \Gamma_i \setminus \{e_i\}$ for $i \in \mathcal{I}$. Furthermore we write $\Gamma_\infty = V_\infty$ and $\Gamma^\times := \Gamma \setminus \{e\}$.

The random walk on Γ is constructed as follows: standing at $x \in \Gamma$ we allow walking to xg with $g \in \bigcup_{i \in \mathcal{I}} \Gamma_i^\times$ in one step. Choose probability measures μ_i on Γ_i^\times for each $i \in \mathcal{I}$ such that μ_i defines an irreducible random walk on Γ_i , that is, $p_i(x, y) = \mu_i(x^{-1}y)$ for all $x, y \in \Gamma_i$. Let $\alpha_1, \dots, \alpha_r > 0$ with $\sum_{i \in \mathcal{I}} \alpha_i = 1$. Then we define the transition probabilities as

$$p(x, xg) := \alpha_i \cdot \mu_i(g)$$

for all $x \in \Gamma$, $g \in \Gamma_i^\times$, and we set $p(x, y) := 0$ otherwise. As the transition probabilities depend only on the increment $g \in \Gamma_i^\times$ we write $\mu(g) := p(x, xg)$ for all $g \in \bigcup_{i \in \mathcal{I}} \Gamma_i^\times$, and $\mu(g) := 0$ otherwise. Analogously, the n -step transition probabilities are given by the convolution powers $\mu^{(n)}$ of μ .

By Theorem 2.5 the random walk converges \mathbb{P}_e -a.s. to a random variable Z_∞ with values in Γ_∞ . Denote by ν the distribution of Z_∞ . Let

$$E_i = \{x_1 x_2 \cdots \in \Gamma_\infty \mid \tau(x_1) = i\} \text{ for } i \in \mathcal{I}.$$

Then ν is uniquely determined by its values on the Borel sets B of the form $x E_i = \{xh \mid h \in E_i\}$ with $i \in \mathcal{I}$, $x \in \Gamma$ and $\tau(x) \neq i$. We will now give a formula for these values:

LEMMA 5.1. *Let $i \in \mathcal{I}$, $x \in \Gamma$ with $\tau(x) \neq i$. Then*

$$\nu(x E_i) = \mathbb{P}_e[Z_\infty \in x E_i] = F(e, x|1)(1 - (1 - \xi_i) G_i(e_i, e_i|\xi_i)).$$

PROOF. The proof of this lemma is extrapolated from Woess [21, Theorem 4 c], where one can find an incorrect formula, which we correct here. First we have

$$\nu(x E_i) = F(e, x|1) \cdot \nu(E_i).$$

Recall that we have by vertex-transitivity $G_i(o_i, o_i|z) = G_i(y, y|z)$ for all $i \in \mathcal{I}$ and all $y \in \Gamma_i$. By Lemma 3.5 we obtain

$$\nu(E_i) = G(o, o|1) \cdot \rho(i) = 1 - (1 - \xi_i) G_i(e_i, e_i|\xi_i).$$

This leads to the proposed formula. \square

Now we reformulate our problem for finding a formula for ℓ . For this purpose, we apply a technique which was used by Ledrappier [10, Section 4 b] for free groups.

By Corollary 3.3 and Lebesgue's Dominated Convergence Theorem we have $\mathbb{P}_e - a.s.$

$$\lim_{n \rightarrow \infty} \frac{\mathbb{E}[\ell(Z_n)]}{n} = \lim_{n \rightarrow \infty} \int \frac{\ell(Z_n)}{n} d\mathbb{P}_e = \int \lim_{n \rightarrow \infty} \frac{\ell(Z_n)}{n} d\mathbb{P}_e = \int \ell d\mathbb{P}_e = \ell.$$

Thus it is sufficient to prove convergence of the sequence

$$\left(\mathbb{E}[\ell(Z_{n+1})] - \mathbb{E}[\ell(Z_n)] \right)_{n \in \mathbb{N}}$$

and to compute its limit, which then must equal ℓ . First we have

$$\mathbb{E}[\ell(Z_n)] = \sum_{h \in \Gamma} \ell(h) \mu^{(n)}(h)$$

and

$$\mathbb{E}[\ell(Z_{n+1})] = \sum_{g, h \in \Gamma} \ell(gh) \mu(g) \mu^{(n)}(h).$$

On the other hand,

$$\mathbb{E}[\ell(Z_n)] = \sum_{g \in \Gamma} \mu(g) \mathbb{E}[\ell(Z_n)] = \sum_{g, h \in \Gamma} \mu(g) \ell(h) \mu^{(n)}(h).$$

Thus we obtain

$$\begin{aligned} \mathbb{E}[\ell(Z_{n+1})] - \mathbb{E}[\ell(Z_n)] &= \sum_{g \in \Gamma} \mu(g) \sum_{h \in \Gamma} (\ell(gh) - \ell(h)) \mu^{(n)}(h) \\ &= \sum_{g \in \Gamma} \mu(g) \int_{\Gamma} (\ell(gZ_n) - \ell(Z_n)) d\mathbb{P}_e. \end{aligned}$$

Define now the random variables

$$Y_n := \ell(gZ_n) - \ell(Z_n)$$

for any given $g \in \bigcup_{i \in \mathcal{I}} \Gamma_i^\times$. We have $Y_n \in \{-1, 0, 1\}$ for all $n \in \mathbb{N}$. By vertex-transitivity gZ_n converges to gZ_∞ . Hence, Y_n converges to a random variable Y_∞ with values in $\{-1, 0, 1\}$ depending only on g and the first block of Z_∞ . In other words, Y_n becomes constant, if n is big enough. If $Z_\infty = x_1 x_2 \dots$, we obtain for given $g \in \bigcup_{i \in \mathcal{I}} \Gamma_i^\times$:

$$Y_\infty = \begin{cases} 0 & , \text{ if } \tau(x_1) = \tau(g) \text{ and } x_1 g \neq e \\ -1 & , \text{ if } \tau(x_1) = \tau(g) \text{ and } x_1 g = e \\ 1 & , \text{ if } \tau(x_1) \neq \tau(g) \end{cases}$$

By Lebesgue's Dominated Convergence Theorem, we infer that

$$\int (\ell(gZ_n) - \ell(Z_n)) d\mathbb{P}_e \xrightarrow{n \rightarrow \infty} \int Y_\infty d\mathbb{P}_e.$$

Consider the function

$$f : \left(\bigcup_{i \in \mathcal{I}} \Gamma_i^\times \right) \times \Gamma_\infty \rightarrow \{-1, 0, 1\} : (g, x_1 x_2 \dots) \mapsto \ell(gx_1) - \ell(x_1)$$

and its projections $f_g : \Gamma_\infty \rightarrow \{-1, 0, 1\} : w \mapsto f(g, w)$ for every $g \in \bigcup_{i \in \mathcal{I}} \Gamma_i^\times$. Observe that each f_g is measurable and thus

$$\int_{\Gamma_\infty} Y_\infty d\mathbb{P}_e = \int_{\Gamma_\infty} f(g, Z_\infty) d\nu = \int_{\Gamma_\infty} f_g(w) d\nu(w).$$

Denote by E_h the event that Z_∞ has as first block the element $h \in \bigcup_{i \in \mathcal{I}} \Gamma_i^\times$ and denote $E_{\neq i}$ the event that Z_∞ starts with a block element not of type $i \in \mathcal{I}$. Then we obtain for $g \in \Gamma_i$

$$\nu(E_{g^{-1}}) = F(e, g^{-1}|1) \cdot (1 - \nu(E_i)) = F(e, g^{-1}|1) \cdot (1 - \xi_i) \cdot G_i(e_i, e_i|\xi_i)$$

and

$$\nu(E_{\neq i}) = (1 - \xi_i) \cdot G_i(e_i, e_i | \xi_i).$$

Writing $G_i(z)$ for $G_i(e_i, e_i | z)$, as before, we can conclude:

$$\begin{aligned} & \mathbb{E}[\ell(Z_{n+1})] - \mathbb{E}[\ell(Z_n)] \\ & \xrightarrow{n \rightarrow \infty} \sum_{g \in \bigcup_{i \in \mathcal{I}} \Gamma_i^\times} \mu(g) \int_{\Gamma^\infty} f_g(w) d\nu(w) \\ & = \sum_{g \in \bigcup_{i \in \mathcal{I}} \Gamma_i^\times} \mu(g) (-\nu(E_{g^{-1}}) + \nu(E_{\neq i})) \\ & = \sum_{i \in \mathcal{I}} \alpha_i (1 - \xi_i) G_i(\xi_i) \left(1 - \underbrace{\sum_{g \in \Gamma_i^\times} \mu_i(g) F_i(e_i, g^{-1} | \xi_i)}_{= \frac{G_i(\xi_i) - 1}{\xi_i \cdot G_i(\xi_i)}}\right) \\ & = \sum_{i \in \mathcal{I}} \alpha_i \frac{1 - \xi_i}{\xi_i} (1 - (1 - \xi_i) G_i(\xi_i)). \end{aligned}$$

Thus we get the rate of escape of the block length as

$$(9) \quad \ell = \sum_{i \in \mathcal{I}} \alpha_i \frac{1 - \xi_i}{\xi_i} (1 - (1 - \xi_i) G_i(\xi_i)).$$

Observe that the technique presented in this section can be extended to a free product of an infinite, countable number of groups. All the required properties of the generating functions used also hold in this case. Furthermore, Y_n is again bounded such that finiteness of ℓ is ensured. Thus the same computations prove the same formula for ℓ if $\mathcal{I} = \mathbb{N}$.

6. Examples

We present two sample applications of our formulas for the rate of escape of the block length. First we look at a free product arising from non-Cayley graphs and then we look at a free product of infinite groups. Note that both examples go beyond previously investigated graph structures for the computation of ℓ .

6.1. Free Product arising from Non-Cayley-Graphs Consider the sets $V_1 = \{A, B, C, D, E, F, o_1\}$, $V_2 = \{G, H, o_2\}$ and $V_3 = \{I, J, o_3\}$ and the random walks on these sets. Their transition probabilities are sketched in Figure 1. Note that none of the graphs in this figure is a Cayley graph. Consider now the corresponding random walk on the free product $V = V_1 * V_2 * V_3$, where $\alpha_1 = 5/9$ and $\alpha_2 = \alpha_3 = 2/9$.

We obtain the following generating functions:

$$\begin{aligned} U_1(o_1, o_1 | z) &= \frac{3}{5}z^2 + \frac{2}{5}z^3, \\ G_1(o_1, o_1 | z) &= \frac{1}{1 - U_1(o_1, o_1 | z)} = \frac{1}{1 - \frac{3}{5}z^2 - \frac{2}{5}z^3}, \end{aligned}$$




FIGURE 1. Some non-Cayley graphs

$$\begin{aligned}
F_1(A, o_1|z) &= F_1(E, o_1|z) = \frac{1}{2}z + \frac{1}{2}z^2, \\
F_1(C, o_1|z) &= z^2, \quad F_1(D, o_1|z) = F_1(F, o_1|z) = z, \\
G_2(o_2, o_2|z) &= \frac{1}{1 - z^2}, \\
F_2(G, o_2, z) &= F_2(H, o_2|z) = z, \\
\bar{H}_1(z) &= \frac{4}{9}z\xi_2(z), \\
\bar{H}_2(z) &= \bar{H}_3(z) = \frac{2}{9}z\xi_2(z) + \frac{5}{9} \cdot \frac{1}{5}z(3\xi_1(z) + 2\xi_1(z)^2).
\end{aligned}$$

Note that $\bar{H}_2(z) = \bar{H}_3(z)$ follows by symmetry. The Green function $G(o, o|z)$ of the corresponding random walk on V has radius of convergence greater than 1. This can be shown by constructing recursive equations using $\bar{H}_1(z)$ and $\bar{H}_2(z)$ and numerical evaluation. Consider

$$\begin{aligned}
\xi_1(z) &= \frac{\frac{5}{9}z}{1 - \bar{H}_1(z)} = \frac{\frac{5}{9}z}{1 - \frac{4}{9}z\xi_2(z)} \quad \text{and} \\
\xi_2(z) &= \xi_3(z) = \frac{\frac{2}{9}z}{1 - \bar{H}_2(z)} = \frac{\frac{2}{9}z}{1 - \frac{2}{9}z\xi_2(z) - \frac{1}{9}z(3\xi_1(z) + 2\xi_1(z)^2)}.
\end{aligned}$$

Substituting $\xi_1(z)$ into $\xi_2(z)$ we have to solve an equation in the variable $\xi_2(z)$. Solving this equation with MATHEMATICA we obtain four continuous solutions, but only one solution satisfies $\xi_2(1) < 1$. Hence, we get $\xi_2(z)$ as this solution and obtain $\xi_1(z)$ from this $\xi_2(z)$. We find that

$$\xi_1(1) \approx 0.66571 \quad \text{and} \quad \xi_2(1) = \xi_3(1) \approx 0.37231.$$

We compute ℓ using Theorem 3.3. The transition matrix of the Markov chain $(\tau_k)_{k \in \mathbb{N}}$ of the alternating vertex types is

$$(\hat{q}(i, j))_{1 \leq i, j \leq 3} = \begin{pmatrix} 0 & 0.5 & 0.5 \\ 0.62769 & 0 & 0.37231 \\ 0.62769 & 0.37231 & 0 \end{pmatrix}$$

and from this we obtain the corresponding invariant probability measure ν with

$$\nu(1) = 0.38563 \quad \text{and} \quad \nu(2) = \nu(3) = 0.30718.$$

Now we are able to compute the rate of escape of the block length to the random walk on V . We obtain

$$\ell \approx 0.33089.$$

If ℓ is computed by Equation (8), then the numerical approximated result and the above result coincide in the first 50 decimal numbers. So numerical approximations do not lead to a distortion of the result.

6.2. $\mathbb{Z}^2 * \mathbb{Z}/2$ Consider $V_1 = \mathbb{Z}^2$ and the simple random walk on it given by $\mu_1((\pm 1, 0)) = \mu_1((0, \pm 1)) = 1/4$. Also, consider the group $V_2 = \mathbb{Z}/2$ and the simple random walk on it given by $\mu_2(1_2) = 1$. We are interested now in the simple random walk on $V = V_1 * V_2$, where $\alpha_1 = 4/5$ and $\alpha_2 = 1/5$. For the computation of ℓ we use Equation (9). Therefore it is sufficient to compute ξ_1, ξ_2 and $G_1((0, 0), (0, 0)|\xi_1)$. For this purpose, we use the computations and results in Woess [23, pages 100, 105, 109]; compare also with Soardi [18].

Before we can compute these values, we have to introduce some auxiliary functions. In the following let the subindex 0 correspond to the random walk on V . Denote

$$W_i(z) = z \cdot G_i(o, o|z) \text{ for } i \in \{0, 1, 2\}.$$

As $W_i(z)$ is strictly increasing, there is an inverse function $W_i^{-1}(z)$ such that $W_i^{-1}(W_i(z)) = z$ holds. By [23, Theorem 9.10] we have

$$G_i(o, o|z) = \Phi_i(zG_i(o, o|z)), \quad \text{where } \Phi_i(t) = \frac{t}{W_i^{-1}(t)} \quad \forall i \in \{0, 1, 2\}.$$

By [23, Example 9.15 (3)],

$$W_1(z) = \frac{1}{4\pi^2} \int_{(-\pi, \pi]^2} \frac{2z}{2 - z \cdot (\cos x_1 + \cos x_2)} d\underline{x},$$

where $\underline{x} = (x_1, x_2)$. Furthermore, by [23, Theorem 9.19] we have the equation

$$\Phi_0(t) = \Phi_1(\alpha_1 t) + \Phi_2(\alpha_2 t) - 1.$$

By [23, Example 9.15 (1)] we have

$$\Phi_2(t) = \frac{1}{2} \left(\sqrt{1 + 4t^2} + 1 \right).$$

This yields

$$\Phi_1\left(\frac{4}{5}W(z)\right) = G(o, o|z) - \frac{1}{2} \left(\sqrt{1 + \frac{4}{25}W(z)^2} - 1 \right).$$

Inverting this equation and multiplication with $4/5 \cdot W(z)$ leads to

$$\frac{\frac{4}{5}W(z)}{\Phi_1\left(\frac{4}{5}W(z)\right)} = \frac{\frac{4}{5}W(z)}{G(o, o|z) - \frac{1}{2} \left(\sqrt{1 + \frac{4}{25}W(z)^2} - 1 \right)}.$$

Applying W_1 onto both sides of this equation yields

$$\frac{4}{5}W(z) = W_1 \left(\frac{\frac{4}{5}W(z)}{G(o, o|z) - \frac{1}{2} \left(\sqrt{1 + \frac{4}{25}W(z)^2} - 1 \right)} \right).$$




FIGURE 2. Graphs of $W_1(z)$ and $(4-5z)z/(4-10z+6z^2)$

Note that we have

$$(10) \quad W_1(\xi_1(z)) = \frac{4}{5}W(z).$$

and thus

$$\xi_1(z) = \frac{\frac{4}{5}W(z)}{\frac{W(z)}{z} - \left(\frac{1}{2}\sqrt{1 + \frac{4}{25}W(z)^2} - 1\right)}.$$

Substituting $y = 4/5 \cdot W(z)$ we obtain

$$\xi_i = \xi_1(1) = \frac{y}{\frac{5y}{4} - \left(\frac{1}{2}\sqrt{1 + \frac{1}{4}y^2} - 1\right)}$$

or equivalently

$$y = \frac{(4-5\xi_1)\xi_1}{4-10\xi_1+6\xi_1^2}$$

Substituting y into Equation (10), we have to solve

$$\frac{(4-5\xi_1)\xi_1}{4-10\xi_1+6\xi_1^2} = W_1(\xi_1)$$

in the unknown variable ξ_1 . The solution can be computed only numerically. Considering the graphs of the functions $(4-5z)z/(4-10z+6z^2)$ and $W_1(z)$ we see that there is only one possible intersection point greater than $4/5$. See Figure 2. Using the bisection method and numerical integration and evaluation we obtain

$$\xi_1 \approx 0.84426, \text{ and also } W(1) = \frac{5}{4}y \approx 1.40724.$$

This yields

$$G_1((0,0), (0,0)|\xi_1) = \frac{\frac{4}{5}W(1)}{\xi_1(1)} \approx 1.33347.$$

By

$$\xi_1 = \frac{\frac{4}{5}}{1 - \frac{1}{5}\xi_2}.$$

we obtain

$$\xi_2 \approx 0.26212, \text{ and } G_2(0,0|\xi_2) = \frac{1}{1-\xi_2^2} \approx 1.07378.$$

Now we have computed all necessary characteristical numbers and the rate of escape of the block length of the simple random walk on $\mathbb{Z}^2 * \mathbb{Z}/2$ can be computed by Equation (9) as

$$\ell \approx 0.23386.$$

7. Remarks

We can extend our considerations in order to compute other types of typical rates of escapes concerning the random walk on the free product:

- I. For $x = x_1 \dots x_n \in V$ and $i \in \mathcal{I}$ the partial block length of x with respect to V_i is given by

$$\ell_i(x) := |\{j \mid j \in \{1, \dots, n\}, x_j \in V_i\}|.$$

As $\ell_i(W_k)/k$ converges for $k \rightarrow \infty$ to $\nu(i)$, which is the invariant probability measure on \mathcal{I} with respect to the Markov chain $(\tau(W_k))_{k \in \mathbb{N}}$, we obtain the partial rate of escape of the block length

$$\frac{\ell_i(Z_n)}{n} \xrightarrow{n \rightarrow \infty} \nu(i) \cdot \ell \quad \mathbb{P}_o - a.s..$$

- II. The set V carries a *Markovian distance* defined by

$$d(x, y) := \min\{n \in \mathbb{N} \mid p^{(n)}(x, y) > 0\},$$

where $x, y \in V$. Note that in general $d(\cdot, \cdot)$ is not necessarily symmetric. The *Markovian length* is defined as $|x| = d(o, x)$. We can extend the considerations of Section 3 for the proof of the existence of the rate of escape of the Markovian length, that is

$$\lambda = \lim_{n \rightarrow \infty} \frac{1}{n} |Z_n|.$$

This yields also a formula for λ given by

$$\lambda = \ell \cdot \sigma, \text{ where } \sigma = \sum_{\substack{i, j \in \mathcal{I}, \\ i \neq j}} \nu(i) \frac{\alpha_j}{\alpha_i} \frac{\xi_i}{\xi_j} \frac{1 - \xi_j}{1 - \xi_i} \sum_{m \geq 1} m \cdot \sum_{y \in S_j(m)} L_j(o_j, y | \xi_j)$$

and $S_i(m) = \{x \in V_i \mid |x| = m\}$ for $i \in \mathcal{I}$ and $m \in \mathbb{N}$.

Acknowledgement

The author is grateful to Wolfgang Woess for numerous discussions on several problems and his help during the preparation of this article, and also to Donald Cartwright for several hints regarding content and exposition.

References

- [1] D. Cartwright, V. Kaimanovich, and W. Woess. Random walks on the affine group of local fields and of homogenous trees. *Ann. Inst. Fourier (Grenoble)*, 44:1243–1288, 1994.

- [2] D. Cartwright and P. Soardi. Random walks on free products, quotients, and amalgams. *Nagoya Math. J.*, 102:163–180, 1986.
- [3] Y. Derriennic. Quelques applications du théorème ergodique sous-additif. *Astérisque*, 74:183–201, 1980.
- [4] A. Dyubina. Characteristics of random walks on wreath products of groups. *J. of Math. Sciences*, 107(5):4166–4171, 2001.
- [5] A. Erschler. On the asymptotics of drift. *J. of Math. Sciences*, 121(3):2437–2440, 2004.
- [6] H. Furstenberg. Non commuting random products. *Trans. Amer. Math. Soc.*, 108:377–428, 1963.
- [7] Y. Guivarc’h. Sur la loi des grands nombres et le rayon spectral d’une marche aléatoire. *Astérisque*, 74:47–98, 1980.
- [8] V. Kaimanovich and A. Vershik. Random walks on discrete groups: boundary and entropy. *Ann. Probab.*, 11:457–490, 1983.
- [9] J. Kingman. The ergodic theory of subadditive processes. *J. Royal Stat. Soc., Ser. B*, 30:499–510, 1968.
- [10] F. Ledrappier. Some asymptotic properties of random walks on free groups. In *CRM Proceedings and Lecture Notes*, volume 28, pages 117–152. CRM, 2001.
- [11] R. Lyons, R. Pemantle, and Y. Peres. Random walks on the lamplighter group. *Ann. of Probability*, 24(4):1993–2006, 1996.
- [12] J. Mairesse. Random walks on groups and monoids with a markovian harmonic measure. *Electron. J. Probab.*, 10:1417–1441, 2005.
- [13] J. Mairesse. Randomly growing braid on three strands and the manta ray. *Ann. Appl. Probab.*, 17(2):502–536, 2007.
- [14] J. Mairesse and F. Mathéus. Random walks on free products of cyclic groups. *J. London Math. Soc.*, 75(1):47–66, 2007.
- [15] J. McLaughlin. *Random walks and convolution operators on free products*. PhD thesis, New York Univ., 1986.
- [16] T. Nagnibeda and W. Woess. Random walks on trees with finitely many cone types. *J. Theoret. Probab.*, 15:399–438, 2002.
- [17] S. Sawyer and T. Steger. The rate of escape for anisotropic random walks in a tree. *Probab. Theory Related Fields*, 76:207–230, 1987.
- [18] P. Soardi. Simple random walks on $\mathbb{Z}^2 * \mathbb{Z}_2$. *Symposia Math.*, 29:303–309, 1986.
- [19] N. T. Varopoulos. Long range estimates for markov chains. *Bull. Sc. math.*, 109:225–252, 1985.

- [20] D. Voiculescu. Addition of certain non-commuting random variables. *J. Funct. Anal.*, 66:323–346, 1986.
- [21] W. Woess. A description of the martin boundary for nearest neighbour random walks on free products. *Probability Measures on Groups*, VII:203–215, 1985.
- [22] W. Woess. Nearest neighbour random walks on free products of discrete groups. *Boll. Un. Mat. Ital.*, 5-B:961–982, 1986.
- [23] W. Woess. *Random Walks on Infinite Graphs and Groups*. Cambridge University Press, 2000.

University of Technology Graz, Institut für Mathematik C, Steyrergasse 30,
A-8010 Graz, Austria
e-mail: gilch@TUGraz.at
URL: <http://www.math.tugraz.at/~gilch/>